

Table I. *Clinical history of the patients with non-organic dyspepsia (NOD) vs. other causes of abdominal pain. TP: True positive; FN: false negative; FP: false positive; TN: true negative.*

Clinical history variable	Positive endpoint	Negative endpoint	TP	FN	FP	TN
1. Location of initial pain	Right or left upper abdomen	Other	39	11	452	831
2. Location of pain at diagnosis	Right or left upper abdomen	Other	37	13	425	858
3. Duration of pain: Duration of pain at diagnosis	>12 hours	≤12 hours	42	8	830	453
4. Intensity of abdominal pain	Subjectively moderate or weak pain	Intolerable pain	42	8	1,075	208
5. Progression of pain from onset to diagnosis	Weaker or subjectively same pain than at the onset	Worse pain	33	17	913	370
6. Type of pain	Steady pain	Colicky or intermitted pain	27	23	704	579
7. Aggravating factors	Movement, coughing, respiration, food or other	No aggravating factors	14	36	342	941
8. Relieving factors	Yes	No	37	13	862	421
9. Previous similar pain	Yes	No	33	17	414	855
10. Vertigo	No	Yes	47	2	1,242	38
11. Nausea	Yes	No	19	31	548	735
12. Vomiting	Yes	No	30	20	545	738
13. Appetite	No appetite	Normal appetite	43	7	934	349
14. Previous indigestion	Yes	No	22	28	257	1,024
15. Jaundice	No	Yes	47	3	1,253	30
16. Bowels	Diarrhea, constipation, blood, mucus or white stools	Normal	16	34	302	981
17. Micturition	Normal	Abnormal	48	2	1,199	84
18. Drugs for abdominal pain	Yes	No	8	42	46	1,236
19. Previous abdominal surgery	Yes	No	28	22	305	977
20. Previous abdominal diseases	Yes	No	22	28	211	1,070
21. Use of alcohol	No	Yes	47	3	1,218	64
22. Gender	Male	Female	32	18	604	679