

Table V. Univariate and multivariate logistic analyses of significant clinicopathological factors predictive of lymphocyte-predominant breast cancer.

Characteristic		Univariate logistic analysis			Multivariate logistic analysis		
		OR	95% CI	<i>p</i> -Value	OR	95% CI	<i>p</i> -Value
Clinical T-stage	T3-T4 <i>vs.</i> T1-2	0.83	0.24-2.59	0.76	4.91	0.40-59.88	0.18
Clinical N-status	Positive <i>vs.</i> negative	0.68	0.23-1.98	0.47	0.54	0.09-3.20	0.49
Nuclear grade	3 <i>vs.</i> 1-2	1.89	0.59-7.36	0.30	0.62	0.07-5.64	0.67
Ki67	≥20% <i>vs.</i> <20%	2.38	0.38-46.14	0.39	1.19	0.03-53.83	0.93
ER	Negative <i>vs.</i> positive	2.71	0.92-8.24	0.07	0.86	0.11-6.45	0.88
HER2	Positive <i>vs.</i> negative	1.37	0.46-4.02	0.57	1.13	0.15-8.53	0.90
TIL-US score	≥4 <i>vs.</i> <3	52.21	9.43-983.82	<0.001	2.90e-6	0	0.71
TIL-CEUS score	≥6 <i>vs.</i> <5	104.12	17.78-2012.52	<0.001	6.61e7	0	0.001

CEUS: Contrast-enhanced ultrasonography; CI: confidence interval; ER: estrogen receptor; HER2: human epidermal growth factor receptor 2; OR: odds ratio; TIL: tumour-infiltrating lymphocyte; US: ultrasonography. Statistically significant *p*-values are shown in bold.