

Table V. *Univariate and multivariate analyses of risk factors for outlet obstruction.*

Variable	Event rate %	Univariate analysis			Multivariate analysis		
		OR	95% CI	<i>p</i> -Value	OR	95% CI	<i>p</i> -Value
Age \geq 75 years	6 (21.4)	0.486	0.143-1.446	0.232			
Male gender	22 (78.6)	1.504	0.495-5.193	0.616			
BMI \geq 25 kg/m ²	5 (17.9)	0.828	0.210-2.767	>0.99			
Diabetes	4 (14.3)	0.437	0.098-1.507	0.198	0.301	0.079-1.150	0.080
Anastomotic leakage	9 (32.1)	4.320	1.247-15.69	0.012	4.160	1.270-13.60	0.018
Abdominal wall thickness \geq 32.5 mm	5 (17.9)	0.706	0.182-2.314	0.602			
Subcutaneous fat thickness \geq 25.6 mm	2 (7.1)	0.352	0.036-1.732	0.223			
Rectus abdominis thickness \geq 10 mm	14 (50)	4.074	1.448-11.78	0.005	4.350	1.580-12.00	0.005

BMI: Body mass index; CI: confidence interval; OR: odds ratio. Statistically significant *p*-values are shown in bold.