

Table II. *Histopathological features of this case and 12 cases with undifferentiated pleomorphic sarcoma previously reported in the literature.*

Case	Study (year) (ref. number)	Histological tumor depth of involvement	Mitotic activity per high power field	Ki-67	Vimentin	Smooth muscle actin	CD 68	S-100 protein	HMB45	CD34
1	Delgado-Partida (1972) (9)	Sclera and ciliary body	NA	NA	NA	NA	NA	NA	NA	NA
2	Urdiales-Viedma (1983) (10)	NA	NA	NA	NA	NA	NA	NA	NA	NA
3	Margo (1989) (11)	Sclera and ciliary body	NA	NA	NA	NA	NA	NA	NA	NA
4	Pe'er (1990) (12)	Cornea	Abnormal mitotic figures	NA	+	NA	NA	-	NA	NA
5	Pe'er (1990) (12)	Fornix and upper eyelid	Many	NA	+	NA	NA	-	NA	NA
6	Balestrazzi (1991) (13)	Orbital invasion	NA	NA	NA	NA	NA	NA	NA	NA
7	Allaire (1999) (14)	Deeper layers and corneal margins	Intense	NA	+(strong)	+	0,3	-	-	-
8	Kim (2006) (15)	Stroma	4-5	NA	+	+	+	-	-	-
9	Kim (2006) (15)	Stroma	<1	NA	NA	NA	+	-	NA	NA
10	Arora (2006) (16)	Tenon's layer	NA	NA	+	-	-	-	-	NA
11	Milman (2007) (17)	NA	Numerous	NA	+(diffuse)	+(focal)	+(focal)	-	-	-
12	Boehlke (2007) (18)	Lateral rectus muscle sheath	Frequent	+(in a few cells)	+(strong)	+(in some cells)	+(strong)	NA	NA	+(strong)
	This case	Stroma	2,5	+(70~80%)	+	-	-	-	-	-

NA: Not available; +: positive.