

Table IV. *Changes in body weight by pretreatment body weight.*

BW group	No. of patients	BW: median (range)	<i>p</i> -Value*
Less than 40 kg			
Pretreatment BW	6	36.6 (33.9-38.8)	
BW at 12 weeks	6	36.5 (34.8-41.3)	0.4004
BW at 24 weeks	6	34.7 (31.0-41.5)	0.7150
BW at 12 months	4	35.0 (31.2-36.6)	0.7150
40 kg-less than 50 kg			
Pretreatment BW	19	43.0 (40.3-49.6)	
BW at 12 weeks	19	43.2 (36.0-51.0)	0.1023
BW at 24 weeks	17	43.2 (35.0-66.0)	0.7332
BW at 12 months	11	42.8 (35.0-48.5)	0.0125
50 kg-less than 60 kg			
Pretreatment BW	21	54.3 (50.0-59.9)	
BW at 12 weeks	21	53.1 (46.0-59.5)	0.0102
BW at 24 weeks	19	53.3 (45.0-63.0)	0.1639
BW at 12 months	13	50.1 (45.0-56.1)	0.0280
60 kg or more			
Pretreatment BW	17	65.2 (60.0-80.0)	
BW at 12 weeks	17	63.6 (59.2-80.5)	0.0105
BW at 24 weeks	13	64.6 (43.0-81.0)	0.0229
BW at 12 months	9	63.1 (54.7-80.5)	0.3270

BW: Body weight. *Comparison with pretreatment weight of each patient (Wilcoxon signed-rank test).